

Phospholipid-TGT

For Laboratory Use Only

Available as **0.25** or **0.5** mM in vials of 3 mL.

Package unit: 10 x 3 mL

Composition

Phospholipid emulsion containing a mixture of highly purified phosphatidyl choline, phosphatidyl serine and sphingomyelin.

Medium: 0.05 mol/L Tris-HCl, pH 7.6.

Properties

A stabilized phospholipid emulsion with high procoagulant activity.
Swirl gently before use.

Stability

A sealed vial is stable at 2-8°C until the expiry date printed on the label. An opened vial is stable for two weeks at 2-8°C.

Do not freeze.

Application

This phospholipid emulsion is suitable for use in hemostasis research related to procoagulant and anticoagulant pathways such as in thrombin generation (TGT).

The 0.5 mM emulsion is specially suitable for use in TGT assays in the presence of activated Protein C (APC).

Phospholipid 0.25 mM

For Laboratory Use Only

Package unit: 10 x 3 mL

Composition

Phospholipid emulsion containing a mixture of highly purified phosphatidyl choline, phosphatidyl serine and sphingomyelin.

Medium: 0.05 mol/L Tris-HCl, pH 7.6.

Properties

A stabilized phospholipid emulsion with high procoagulant activity.
Swirl gently before use.

Stability

A sealed vial is stable at 2-8°C until the expiry date printed on the label. An opened vial is stable for one month at 2-8°C.

Do not freeze.

Application

This phospholipid emulsion is suitable for use in hemostasis research related to procoagulant and anticoagulant pathways.