



## Goat anti-human Protein C (PC)

Affinity-Purified IgG

0.5 mg

**Product #:** GAPC-AP

**Lot #:** XXXX

**Expiry date:** XXXX

Store at -10 to -20°C

For Research Use Only.

Not for use in diagnostic procedures.

### Description of Protein C (PC)

Protein C (PC) is a vitamin K-dependent glycoprotein produced in the liver. The concentration of PC in plasma is ~4 µg/ml (~60 nM). A deficiency of Protein C (quantitative or qualitative) is a risk factor for vascular thrombosis. Protein C is expressed as a two-chain molecule with a molecular weight of 62 kDa. The light chain (21 kDa) of PC consists of two EGF-like domains and an amino-terminal domain containing one hydroxyaspartic acid and 11 γ-carboxyglutamic acid (gla) residues. These residues allow PC to bind to membranes that contain acidic phospholipids in a calcium dependent manner. The heavy chain of PC (41 kDa) consists of the catalytic domain and an activation peptide. Activation of Protein C results from cleavage at residue Arg<sup>12</sup> in the heavy chain by a complex of thrombin and a cell surface cofactor thrombomodulin. The activation of PC is associated with the release of a small activation peptide (2-3 kDa, called Protein C peptide, or PCP) from the N-terminal of the heavy chain. Activated Protein C (APC) is a serine protease with anticoagulant activity. APC, in complex with a phospholipid membrane, calcium and the Protein S cofactor, exhibits anticoagulant activity through the proteolytic inactivation of coagulation cofactors Va and VIIIa. The primary inhibitor of APC activity in plasma is Protein C Inhibitor (PCI, also called Plasminogen Activator Inhibitor-3, PAI-3) and to a lesser extent by α<sub>1</sub>antitrypsin and α<sub>2</sub>macroglobulin. The inhibitory activity of PCI is stimulated approximately 10 fold by heparin<sup>1-3</sup>.

### REFERENCES and REVIEWS

1. Broze GJ, Miletich JP; Biochemistry and Physiology of Protein C, Protein S and Thrombomodulin; in Hemostasis and Thrombosis, 3<sup>rd</sup> Edition, eds. RW Colman, J Hirsh, VJ Marder and EW Salzman, pp 259-276, J.B. Lippincott Co., Philadelphia PA, USA, 1994.
2. Esmon CT, Esmon NL, Le Bonniec B, Johnson AE; Protein C Activation; Methods in Enzymology 222, pp 359-385, 1993.
3. Heeb MJ, Mosher D, Griffin JH; Activation and Complexation of Protein C and Cleavage and Decrease of Protein S in Plasma of Patients With Intravascular Coagulation; Blood 73, pp 455-461, 1989.

### Product Specifications

#### Description:

Vial containing XXXX ml of IgG purified by affinity-chromatography on immobilized PC. Total protein is 0.5 mg.

#### Format:

Affinity-purified IgG (APIgG), clear liquid.

#### Host Animal:

Goat

#### Immunogen:

Human Protein C purified from plasma.

#### Concentration:

APIgG concentration is XXXX mg/ml, determined by absorbance using an extinction coefficient ( $E_{280}^{1\%}$ ) of 13.4.

#### Buffer:

10 mM HEPES, pH 7.4, 150 mM NaCl, 50% (v/v) glycerol.

#### Storage:

Store between -10 and -20°C. Product will become viscous but will not freeze. Avoid storage in frost-free freezers. Keep vial tightly capped. Allow product to warm to room temperature and gently mix before use.

#### Specificity:

This antibody is specific for Protein C as demonstrated by immunoelectrophoresis and ELISA.

#### Applications:

Suitable as a source of enriched antibodies to human PC.

#### Neutralizing activity:

Not determined.

#### Species Cross Reactivity:(immunodiffusion vs. citrated plasma)

Human:	XXXX	Mouse:	XXXX	Rat:	XXXX
Rabbit:	XXXX	Pig:	XXXX	Dog:	XXXX