

Porcine Fibrinogen

100 mg

Ref#: PFG
Lot#: xxxxxx
Exp. Date: xxxx-xx



For Research Use Only
Not for Use in Diagnostic Procedures
For *in vitro* use only

Description:	Fibrinogen (Porcine)
Format:	Lyophilized in 20 mM sodium citrate-HCl / pH 7.3
Host:	Porcine
Storage:	Store between +2 and +8°C After reconstitution aliquot and freeze at ≤-60°C
Volume:	1 vial containing 4.845 mL
Total Protein:	100 mg
Concentration:	20.64 mg/mL by Absorbance; Extinction Coefficient $E^{1\%_{280}} = 15.1$
Activity:	96.90% Clottable
Molecular weight:	330.000 daltons

Fibrinogen (or Factor I) is an acute phase protein that is part of the coagulation cascade of proteins. Thrombin rapidly proteolyzes fibrinogen, releasing fibrinopeptide A. The loss of this small peptide is not sufficient to make the resulting fibrin molecule insoluble, but it tends to form complexes with adjacent fibrin and fibrinogen molecules. Thrombin then cleaves a second peptide, fibrinopeptide B, from fibrin and the fibrin monomers formed then polymerize spontaneously to form an insoluble gel. The polymerized fibrin is held together by noncovalent and electrostatic forces and stabilized by the transamidating enzyme, factor XIIIa, that is produced by the action of thrombin on factor XIII. The insoluble fibrin aggregates (clots) and aggregated platelets then block the damaged blood vessel and prevent further bleeding. The amount of fibrinogen in the plasma can serve as a nonspecific indicator of whether or not an inflammatory process is present in the body. Fibrinogen from any mammalian source will be cleaved by thrombin from any mammalian source.

The Porcine Fibrinogen was purified from fresh frozen porcine plasma using a combination of salt precipitations and column chromatography. The purified protein is homogeneous on SDS-PAGE and shows the above mentioned clottability in a thrombin based assay.

For Research Use Only. Not for Human Use.

The above protein was purified from Human plasma that was tested and found negative by FDA accepted methods for Anti-HIV 1/2, Anti-HTLV I & II, HBsAg, Anti-HCV, Syphilis, HBC Ab, HIV-1 p24 Ag or HIV-1 RNA, HCV RNA and HBV RNA. Donors are screened for CJD (Creutzfeld-Jakob Disease).